



EVALUATION SALIVARY LEVELS OF LACTOPEROXIDASE AND LACTOFERRIN IN PRESCHOOL CHILDREN WITH DENTAL CARIES

Raghda Lutfi Nussrat^{1*}, Zainab A. Aldhaher²

¹Department of Basic science, College of Dentistry, University of Baghdad, Iraq, raghda.lotfi2207m@codental.uobaghdad.edu.iq

²Professor, Department of Basic Science, College of Dentistry, University of Baghdad, Iraq, zainab.aldhahir@codental.uobaghdad.edu.iq

Received 29/ 11/ 2023, Accepted 21/ 1/ 2024, Published 31/ 12/ 2025



This work is licensed under a CCBY 4.0 <https://creativecommons.org/licenses/by/4.0>

ABSTRACT

This study was conducted to determine the saliva levels of lactoperoxidase and lactoferrin between children who do not suffer from tooth decay and those who have early childhood caries. Eighty healthy children of kindergarten age (aged between 4-5 years) were enrolled in this study, according to the World Health Organization, in the city of Baghdad. An oral examination was performed and 40 kindergarten children were suffering from severe caries, which was detected. According to Wyne classification, it was considered a (study group), while 40 caries-free kindergarten children were considered a (control group) that matched by age, sex, and socioeconomic status. Salivary lactoperoxidase and lactoferrin levels were assessed using an ELISA device. The results of the statistical analysis showed that the levels of salivary lactoperoxidase and salivary lactoferrin increased significantly in the study group (those with caries) compared to the caries-free group (control group). The research found that lactoperoxidase and lactoferrin can be used as markers to predict the risk of tooth decay in children's saliva.

Keywords: Dental caries, Lactoperoxidase, Lactoferrin, Childhood, Saliva.

تقييم مستويات اللعاب من لاكتوبيروكسيديز واللاكتوفيرين في الأطفال في سن ما قبل المدرسة الذين لديهم تسوس الأسنان

رغدة لطفي نصرت¹, زينب عبد الجبار الظاهر²

¹قسم العلوم الأساسية، كلية طب الأسنان، جامعة بغداد، العراق raghda.lotfi2207m@codental.uobaghdad.edu.iq

²أستاذ، قسم العلوم الأساسية، كلية طب الأسنان، جامعة بغداد، العراق zainab.aldhahir@codental.uobaghdad.edu.iq

الخلاصة

أجريت هذه الدراسة لمعرفة مستويات اللعاب من لاكتوبيروكسيديز واللاكتوفيرين بين الأطفال الذين لا يعانون من تسوس الأسنان وأولئك الذين لديهم تسوس في مرحلة الطفولة المبكرة. حيث تم تسجيل في هذه الدراسة ثمانين طفلاً يتمتعون بصحة جيدة بسن رياض الأطفال (تراوح أعمارهم بين 4-5 سنة) حسب منظمة الصحة العالمية، في مدينة بغداد، وتم إجراء الفحص الفموي وكان 40 طفلاً من أطفال الروضة يعانون من حالة تسوس حاد وتم الكشف عنه وفقاً لتصنيف واين وتم اعتبارها (مجموعة دراسة)، بينما كان 40 طفلاً خالياً من التسوس في اطفال الروضة وتم اعتبارها (مجموعة ضابطة) التي تتطابق مع العمر والجنس والحالة الاجتماعية والاقتصادية. تم تقييم مستويات اللاكتوبيروكسيديز اللعابي واللاكتوفيرين باستخدام جهاز الآليزا. حيث أظهر نتائج التحليل الإحصائي أن مستوى اللاكتوبيروكسيديز اللعابي واللاكتوفيرين العابي ارتفع بشكل ملحوظ في مجموعة الدراسة (المصابين بالتسوس) مقارنة بالمجموعة الضابطة من التسوس (المجموعة الضابطة). توصل البحث إلى أنه يمكن استخدام اللاكتوبيروكسيديز واللاكتوفيرين كعلامات للتنبؤ بمخاطر تسوس الأسنان في لعاب الأطفال.

الكلمات المفتاحية: تسوس الأسنان، لاكتوبيروكسيديز، لاكتوفيرين، الأطفال ، اللعاب.

*This article is taken from the first researcher's master's thesis.



INTRODUCTION

Dental caries is a multifactorial, locally damaging, microbial based illness that is mostly irreversible. The host, the microorganisms, and the diet each play a different role in the development and growth of tooth caries (**Najm & Al-Mizraqchi, 2019**). Moreover dental caries is a multifactorial oral illness caused by a glucose induced plaque that is distinguished by the modal demineralization of the tooth's tough tissues (**Pitts et al., 2017**). therefore saliva play a crucial function as indicators in the identification of caries because saliva around both hard and soft tissue of the mouth contains some inorganic as well as organic substances that include specific elements that significantly contribute to decay in the patient's body (**Ahsan, 2019**). Unbalanced cariogenic bacteria cause dental caries (**Chen et al., 2020**). Thus, The predominant cariogenic microorganisms in biofilm and salivary will produce further acids during the process of fermentation (**Al-zahraa & Aldhaher, 2017**). As a result, the acid destroys the enamel layer form by causing loss of minerals within the hard tissues of the tooth (**Chen et al., 2020**).

Early childhood caries is the condition in which a kid under the age of five has a number of primary teeth that are missing, decaying, or filled. Medically, ECC has several of distinctive traits, such as fast decay development which impacts multiple teeth shortly following their appear in the mouth (**Feldens et al., 2010**). The maxillary incisors' labial surface and the lingual and buccal surfaces of the mandibular and maxillary molars' surface are among the dental sites where it typically occurs (**Anil & Anand, 2017**). Salivary biochemical and microbiological changes may have an impact on the happening and riskiness of tooth decay (**Al-Khayoun & Diab, 2015**). In another hand dental caries may be made more likely by poor oral hygiene practices (**Misbah, 2005**). Children's oral and dental health depends on a healthy diet (**Chaloob, & Qasim, 2013**).

Dental decay can be caused by food or substrate in addition to other variables, and the etiology of early childhood caries is complex. The chance of tooth decay increases when sweets are added to milk or juice. In comparison with kids who don't have these dietary habits, kids who drink sugary drinks at least twice at night are more prone to early childhood caries (**Bachtiar et al., 2019**).

The salivary glands in the mouth produce saliva, which is a watery secretion. Saliva is crucial for the development of caries and for preventing it (**Munther, 2020**). It is composed of water, electrolytes, mucus, enzymes, and antibacterial agents. Saliva may be used to diagnose systemic diseases or as indicators of exposure to many toxic or harmful substances due to the existence of hundreds of various components (**Arunkumar et al., 2014**).

Salivary enzymes such as lactoperoxidase are keeping important functions of maintaining dental health. It is considered as necessary for the host's initial line of defense against infections including *Streptococcus mutans*, *Streptococcus sobrinus*, and *Streptococcus sanguinis*. Salivary materials may offer a lot of medical data. Additionally, this salivary proteomics can result in the identification of clinical indications (**Si et al., 2015**). Salivary lactoperoxidase is a type of protein that has distinctive enzyme properties and guards against microbial breakdown of saliva. It functions together with hydrogen peroxide and thiocyanate.

This produces lead to oxidize bacterial sulfhydryl bonds and impede microbial digestion of glucose (**Gornowicz et al., 2014**). The lactoperoxidase enzyme functions as a catalyst in this reaction. hydrogen peroxide (H_2O_2) and thiocyanate ion undergo an oxidation process and that results hypothiocyanate ions which these substances are in charge of eliminating microorganisms (**Jyoti et al., 2009**), One more potential aid dentin affected by



decay, due to its antimicrobial properties, is a combination of the enzymes lactoferrin, and lactoperoxidase are among the various antimicrobial agents associated with the immune function of saliva (**Jyoti et al., 2009**).

The lactoferrin is non-enzymatic microbial protein. It has been found to be extensively distributed in bodily fluids including saliva, tears, and even white blood cells. The main and minor salivary glands' serous cells release it. The microbes are deprived of this protein crucial due to iron-chelating his activity. Lactoferrin, also known as apolactoferrin when it is iron-free, has the ability to act as an antimicrobial substance by directly interacting with organisms and agglutinating *S. mutans*. This makes it simple to remove the agglutinated organisms from the mouth using the physical forces of saliva and gulping the agglutinated microbes. Additionally, a wide range of types have shown lactoferrin to have significant antimicrobial and antiviral action (**Aziz, 2023**). These substances may have a bactericidal or potentially inhibiting impact on dental infections (**Shimada et al., 2008**). The aim of the investigation was to compare the salivary levels of lactoferrin and lactoperoxidase between children without dental decay and those who had early childhood caries.

MATERIALS AND METHODS

This case control study was conducted from December 2022 to May 2023 in eighty healthy looking Kindergartens children with an age of 4-5 years old according to (**World Health Organization, 2013**), in Baghdad city and oral examination was done and 40 Kindergartens children suffering a severe form of early childhood caries were detected according to (**Wyne, 1999**), class and were considered as study group and 40 Kindergartens caries free children were the control group matching age and gender and socioeconomic status. The Shapiro-Wilk test was run to determine whether the data's distribution was normally distributed.

EXAMINATION OF THE ORAL CAVITY

Oral evaluation was performed using a decay, missing, and filled index in accordance with (**World Health Organization, 2013**). And 40 children suffering a severe form of early childhood caries were selected as study group according to (**Wyne, 1999**).

SALIVA COLLECTION

After the oral examination, make sure the childhood did not eat or drink for at least one hour before saliva collection. Four milliliters of unstimulated saliva were collected by drooling method in plain tube, saliva were obtained collected in the morning between 9-11am. (**Al-Musawi & Ali, 2023**). After this, each tubes was transported in an ice box to the laboratory. To preserve salivary proteins from hydrolysis, saliva was separated in lab from 10 to 15 minutes at (3000) rpm at 28°C. The supernatant was transported to an Eppendorf tube, numbering and freezing under -20°C for identification of markers.

BIOMARKERS DETECTION

The level of lactoperoxidase and lactoferrin were measured as instructed in the brochure. By using a sandwich Enzyme-Linked Immunosorbent Assay (ELISA) kit that is commercially available (Cloud-Clone Corp; USA).



RESULT

There are no significant differences between male and female in age in study and control group as illustrated in table (1).

Table (1): Age distribution of study and control group.

Age	Study group	Control group	p value
	Mean \pm SD	Mean \pm SD	
Male	4.755 ± 0.393	4.435 ± 0.427	0.007*
Female	4.715 ± 0.394	4.615 ± 0.450	0.174 ns
p value	0.762 ns	0.126 ns	

*= significant ns= non-significant

There are no significant differences in gender distribution between study and control group as demonstrated in table (2).

Table (2): Gender distribution of study and control group.

Gender		Control group No=40	Study group No=40	Total
Male	No	20	20	40
	% within group	50	50	50
Female	No	20	20	40
	% within group	50	50	50
Total	No	40	40	80
	% within type	100	100	100




Figure (1): Gender distribution of study and control group.

DETECTION OF SALIVARY LACTOPEROXIDE LEVELS

The results in Tab. (3) and Fig. (2) Revealed increase in salivary lactoperoxidase level in study group (644.6 ± 122) in comparison to control groups (245.9 ± 36.27) having differences that are statistically significant ($P < 0.05$).

Table (3): Descriptive criteria of lactoperoxidase marker of study and control groups.

Descriptive criteria	lactoperoxidase concentration (ng/ml) (n= 40)	
	Control group	Study group
Minimum	137.2	507.4
Maximum	306.8	927.4
Median	231.8	593.2
Range	169.5	419.9
\pm Std. Deviation	36.27	122
Std. Error of Mean	5.735	19.29
P value	$<0.0001^{a*}$	

* Highly significant, a Mann Whitney




Figure (2): Descriptive criteria of lactoperoxidase marker.

DETECTION OF SALIVARY LACTOFERRIN LEVELS

The results in Tab. (4) and Fig. (3) Revealed increase in salivary Lactoferrin level in study group (15.42 ± 2.208) in comparison to control groups (6.812 ± 1.963) having differences that are statistically significant ($P < 0.05$).

Table (4): Descriptive criteria of lactoferrin marker of study groups.

Descriptive criterion	Lactoferrin concentration (ng/ml) (n= 40)	
	Control group	Study group
Minimum	1.831	13
Maximum	9.57	24.85
Median	7.129	15.16
Range	7.739	11.85
±Std. Deviation	1.963	2.208
Std. Error of Mean	0.3103	0.3492
P value	<0.0001 ^{a*}	

* Highly significant, a Mann Whitney




Figure (3): Descriptive criteria of lactoferrin marker.

DISCUSSION

Saliva has a variety of roles in defending the teeth toward abrasion, wear, erosion, and tooth decay (Dawes *et al.*, 2015), where it consists of 99% water, One of the salivary glands' primary jobs is to continuously release saliva into the mouth in order to keep the oral tissue and made it moist and less prone to abrasion as well as to facilitate the removal of microbes , desquamated cells of the epithelium, white blood cell, and food particles during process of swallowing (Dawes *et al.*, 2015; Alobadi, 2020). Several studies have sought in previous years to link certain characteristics of salivary production and content to risk of tooth decay. Furthermore salivary samples are diagnostic tools because saliva is easily collected and analysed, also many of salivary indicators may be able to distinguish between periodontal wellness and illness with accuracy (Abdullameer & Abdulkareem, 2023; Kazem *et al.*, 2023).



Forty children with dental caries (severe type) ECC and forty free caries children participated in the current study and no significant age differences and gender were recorded.

The results also revealed that the study group had higher salivary levels of lactoperoxidase and lactoferrin as compared to the control groups. Regarding lactoperoxidase the findings of this research were in accordance with others someone uncovered that high-intensity caries is linked with increased levels of lactoperoxidase (Gornowicz *et al.*, 2014), another study done by Ruan and *et al.* Also came in agreement with the results of the current study (Ruan *et al.*, 2021). These outcomes might be attributed to the fact that the lactoperoxidase enzyme functions as a catalyst in this reaction. Hydrogen peroxide (H₂O₂) and thiocyanate ion undergo an oxidation process and that results hypothiocyanate ions which these substances are in charge of eliminating microorganisms (Al-Baarri *et al.*, 2011).

The components of the lactoperoxidase structure like lactoperoxidase, hydrogen peroxide, and thiocyanate ions exert an inhibitory effect on cariogenic oral microbiota. The amounts of thiocyanate ions may be raised in vivo by adding more natural enzymes, such as lactoperoxidase, to the saliva and therefore the quantity of cariogenic microbiota in children with Early Childhood Caries shall be decreased by this elevated level of thiocyanate (Jyoti *et al.*, 2009). All these findings support the increased level of lactoperoxidase in saliva of children with dental caries in this study.

In this research also revealed increased level of saliva lactoferrin in children with dental caries these results are supported with the results of researchers who found that lactoferrin was significantly lower in the caries-free group than in the dental caries group (Ruan *et al.*, 2021). Also Felizardo and *et al.* Found Lactoferrin was discovered to be positively linked with Decay Massing and Filling (DMFT) (Felizardo *et al.*, 2010).

Another study noticed was a connection between the DMFT index and lactoferrin expression and declared that dental caries tend to be more likely to arise when lactoferrin is present, additionally a correlation between decay risk /activities and lactoferrin genetic polymorphism discover (Vitorino *et al.*, 2006).

CONCLUSION

Lactoperoxidase and Lactoferrin may be used as markers for dental caries risk prediction in saliva of children.

ACKNOWLEDGMENT

The authors express thanks to the University of Baghdad's College of Dentistry in Iraq for their help in this study.

REFERENCES

1. Abdullameer, M. A., & Abdulkareem, A. A. (2023). Diagnostic potential of salivary interleukin-17, RANKL, and OPG to differentiate between periodontal health and disease and discriminate stable and unstable periodontitis: A case-control study. *Health Science Reports*, 6(2), 11-20.
2. Ahsan, H. (2019). Biomolecules and biomarkers in oral cavity: Bioassays and immunopathology. *Journal of Immunoassay and Immunochemistry*, 40(1), 52-69.



3. Al-Baarri, A. N., Hayashi, M., & Ogawa, M. (2011). Effects of mono-and disaccharides on the antimicrobial activity of bovine lactoperoxidase system. *Journal of Food Protection*, 74(1), 134-139.
4. Al-Khayoun, J. D., & Diab, B. S. (2015). The relation of salivary glucose with dental caries and Mutans Streptococci among type1 diabetic mellitus patients aged 18-22 years. *Journal of Baghdad College of Dentistry*, 27(3), 146-151.
5. Al-Musawi, M. A., & Ali, O. H. (2023). Assessment of salivary interleukin-1 β levels in patients with gingivitis and periodontitis: an analytical cross-sectional study. *Dental Hypotheses*, 14(1), 3-6.
6. Alobadi, E. G. Y. (2020). Study of some immunological parameters in patient that infected with Streptococcus pyogenes. *Iraqi Journal of Market Research and Consumer Protection*, 12(1), 35-43.
7. Al-zahraa, J. J., & Aldhaher, Z. A. (2017). Evaluation of mutans streptococci concerning oral health in the saliva of pregnant women. *Revis Bionatura*, 8 (2) 81-92.
8. Wyne A. (1999). Prevalence and risk factors of nursing caries in Adelaide, South Australia. *international journal of Japanese Society of Pediatric Dentistry*, 9(1), 31-36.
9. Anil, S., & Anand, P. S. (2017). Early childhood caries: prevalence, risk factors, and prevention. *Frontiers in pediatrics*, 5(2), 157-168.
10. Arunkumar, S., Arunkumar, J. S., Krishna, N. B., & Shakunthala, G. K. (2014). Developments in diagnostic applications of saliva in oral and systemic diseases-A comprehensive review. *Journal of Scientific and Innovative Research*, 3(3), 372-387.
11. Aziz, R. A. (2023). Study of the synergistic effect of proteins produced from *Saccharomyces cerevisiae* with lactoferrin against multi resistant diarrheal bacteria. *Iraqi Journal of Market Research and Consumer Protection*, 15(1), 45-53.
12. Bachtiar, Z. A., Primasari, A., & Sungkar, S. (2019). The Effectiveness of Milk Lactoperoxidase in Increasing Salivary Lactoperoxidase Levels in Children with Early Childhood Caries. *Journal of Dental and Medical Sciences*, 3(1), 35-39.
13. Chaloob, E. K. & Qasim, A. A. (2013). Nutritional status in relation to oral health status among patients attending dental hospital. *Journal of Baghdad College of Dentistry*, 25(2) 114-119.
14. Chen, X., Daliri, E. B. M., Kim, N., Kim, J. R., Yoo, D., & Oh, D. H. (2020). Microbial etiology and prevention of dental caries: exploiting natural products to inhibit cariogenic biofilms. *Pathogens*, 9(7), 56-69.
15. Dawes, C. (2012). Salivary clearance and its effects on oral health. *Saliva and oral health*, 5(2), 71-85.
16. Dawes, C., Pedersen, A. L., Villa, A., Ekström, J., Proctor, G. B., Vissink, A., & Wolff, A. (2015). The functions of human saliva: A review sponsored by the World Workshop on Oral Medicine. *Archives of oral biology*, 60(6), 863-874.
17. Feldens, C. A., Giugliani, E. R. J., Duncan, B. B., Drachler, M. D. L., & Vítolo, M. R. (2010). Long-term effectiveness of a nutritional program in reducing early childhood caries: a randomized trial. *Community dentistry and oral epidemiology*, 38(4), 324-332.
18. Felizardo, K. R., Gonçalves, R. B., Schwarcz, W. D., Poli-Frederico, R. C., Maciel, S. M., & Andrade, F. B. D. (2010). An evaluation of the expression profiles of salivary proteins lactoferrin and lysozyme and their association with caries experience and activity. *Revista Odonto Ciência*, 25(2), 344-349.



19. Gornowicz, A., Tokajuk, G., Bielawska, A., Maciorkowska, E., Jabłoński, R., Wójcicka, A., & Bielawski, K. (2014). The assessment of sIgA, histatin-5, and lactoperoxidase levels in saliva of adolescents with dental caries. *International medical journal of experimental and clinical research*, 20(3), 85-95.
20. Jyoti, S., Shashikiran, N., & Subba Reddy, V. (2009). Effect of lactoperoxidase system containing toothpaste on cariogenic bacteria in children with early childhood caries. *Journal of Clinical Pediatric Dentistry*, 33(4), 299-304.
21. Kazem, N. M., Abdulkareem, A. A., & Milward, M. R. (2023). Salivary E-cadherin as a biomarker for diagnosis and predicting grade of periodontitis. *Journal of Periodontal Research*, 6(2), 85-95.
22. Misbah, M. M. (2005). Oral habits in relation to dental caries and gingival health among children attending the dental hospital. *Journal of baghdad college of dentistry*, 17(3), 88-95
23. Munther, S. (2020). The impact of salivary lactoperoxidase and histatin-5 on early childhood caries severity in relation to nutritional status. *The Saudi Dental Journal*, 32(8), 410-416.
24. Najm, A. Z. & Al-Mizraqchi, A. S. (2019). The Dental Caries Experience in Relation to Salivary Flow Rate, sIgA and Mutans Streptococci Bacteria in Smoker and Non-Smoker Patients. *Journal of Baghdad College of Dentistry*, 31(1), 52-58.
25. Pitts, N. B., Zero, D. T., Marsh, P. D., Ekstrand, K., Weintraub, J. A., Ramos-Gomez, F., & Ismail, A. (2017). *Nature reviews Disease primers*, 3(1), 1-16.
26. Ruan, W., Sun, C., Gao, Q., & Shrivastava, N. (2021). Metaproteomics associated with severe early childhood caries highlights the differences in salivary proteins. *Archives of Oral Biology*, 6(3), 105-111.
27. Shimada, J., Moon, S. K., Lee, H. Y., Takeshita, T., Pan, H., Woo, J. I., & Lim, D. J. (2008). Lysozyme M deficiency leads to an increased susceptibility to *Streptococcus pneumoniae*-induced otitis media. *Infectious diseases*, 8(1), 1-11.
28. Si, Y., Ao, S., Wang, W., Chen, F., & Zheng, S. (2015). Magnetic bead-based salivary peptidome profiling analysis for severe early childhood caries. *Caries research*, 49(1), 63-69.
29. Vitorino, R., de Moraes Guedes, S., Ferreira, R., Lobo, M. J. C., Duarte, J., Ferrer-Correia, A. J. & Amado, F. M. (2006). Two-dimensional electrophoresis study of in vitro pellicle formation and dental caries susceptibility. *European journal of oral sciences*, 114(2), 147-153.
30. World Health Organization. (2013). *Oral health surveys: basic methods*. World Health Organization.